

DUAL BRST SYMMETRY FOR QED

R. P. MALIK *

*S. N. Bose National Centre for Basic Sciences,
Block-JD, Sector-III, Salt Lake, Calcutta-700 098, India*

Abstract: We show the existence of a co(dual)-BRST symmetry for the usual BRST invariant Lagrangian density of an Abelian gauge theory in two dimensions of space-time where a $U(1)$ gauge field is coupled to the Noether conserved current (constructed by the Dirac fields). Under this new symmetry, it is the gauge-fixing term that remains invariant and the symmetry transformations on the Dirac fields are analogous to the chiral transformations. This interacting theory is shown to provide a tractable field theoretical model for the Hodge theory. The Hodge dual operation is shown to correspond to a discrete symmetry in the theory and the extended BRST algebra for the generators of the underlying symmetries turns out to be reminiscent of the algebra obeyed by the de Rham cohomology operators of differential geometry.

* E-mail address: malik@bosen.bose.res.in

1 Introduction

The study of symmetries in theoretical physics has brought together many physical ideas and underlying mathematical concepts on an equal footing where both have enriched each other in a grand and illuminating manner. One such symmetry is the Becchi-Rouet-Stora-Tyutin (BRST) symmetry [1,2] which has played a key role in bringing together the physics of gauge theory (endowed with the first-class constraints) and mathematics of differential geometry (related with connections and curvature tensors, etc.) in the realm of what is known as the BRST cohomology. In fact, the nilpotent ($Q_B^2 = 0$) BRST charge (Q_B) is analogous to the exterior derivative d ($d^2 = 0$) of differential geometry and BRST closed states ($Q_B|phys\rangle = 0$) are the *physical states* which are analogous to the closed-forms f ($df = 0$) of differential geometry. The subsidiary condition $Q_B|phys\rangle = 0$ is required to ensure that the operator form of first-class constraints of the original gauge theories must annihilate the physical states of the theory (Dirac's prescription) [3-8]. Now two physical states are said to belong to the same cohomology class with respect to BRST charge Q_B if they differ by a BRST exact state. There are two other operators of differential geometry that are required in the definition of the Hodge decomposition theorem which states that, on a compact manifold, any n -form f_n ($n = 0, 1, 2, \dots$) can be uniquely written as the sum of a harmonic form h_n ($\Delta h_n = 0, dh_n = 0, \delta h_n = 0$), an exact form de_{n-1} and a co-exact form δc_{n+1} as

$$f_n = h_n + d e_{n-1} + \delta c_{n+1}, \quad (1.1)$$

where δ ($\delta = \pm * d *$; $\delta^2 = 0$) is the co(dual)-exterior derivative and Δ ($\Delta = (d + \delta)^2 = d\delta + \delta d$) is the Laplacian operator [7-13]. It has been a long-standing problem to express δ and Δ in the language of some *local* symmetry properties of a given BRST invariant Lagrangian density for a gauge theory. Some very interesting and illuminating attempts have been made towards this goal for the *interacting* (non)Abelian gauge theories in any arbitrary dimension of space-time but symmetry transformations have turned out to be non-local and non-covariant [14-17]. In the covariant formulation [18], the nilpotency is restored only for the specific values of the parameters of the theory.

Recently, it has been shown [19,20] that free two-dimensional (2D) Abelian- and self-interacting non-Abelian gauge theories (without any interaction with matter fields) are the prototype examples of a new kind of topological field theories [†] [21] that provide a field theoretical model for the Hodge theory where all the de Rham cohomology operators (d, δ, Δ) correspond to the *generators* of local, continuous, covariant and nilpotent (for d and δ) symmetries of the theory. Such an analogy has also been established for the free two-form Abelian gauge theory in four ($3 + 1$) dimensional space-time [22]. All these above examples of free gauge theories are, however, ideal as well as simple examples and have very little to do with physical reality. The most natural and physically interesting

[†]A theory with a flat space-time metric but without any propagating degrees of freedom. It captures together some of the key features of Witten- and Schwarz-type of topological field theories.

gauge theories are the interacting theories. It is, therefore, a nice idea to explore the possibility of having an interacting field theory as a tractable model for the Hodge theory. The purpose of the present paper is to show that 2D quantum electrodynamics (QED) (a dynamically closed system of $U(1)$ gauge field A_μ and Dirac fields ψ and $\bar{\psi}$) is an *interacting* field theoretical model for the Hodge theory. Besides the local BRST charge (analogue of d), we show that there are local co-BRST charge Q_D (analogue of δ) and a local bosonic (Casimir) operator W (analogue of Δ) that are generators of symmetry transformations for the BRST invariant Lagrangian density of QED. Like two-form $F = dA$ remains invariant under the BRST transformations and the Dirac fields transform to the analogue of the local gauge transformations, it is the gauge-fixing term † that remains invariant under the dual BRST transformations and the Dirac fields transform as an analogue of the chiral transformations. We show that the existence of a discrete symmetry connects these two symmetry transformations and turns out to be the analogue of the Hodge dual $*$ operation of the differential geometry. Finally, we show that the extended BRST algebra constituted by generators of all the underlying symmetries of the theory mimic the algebra of the de Rham cohomology operators of differential geometry. To the best of our knowledge, this *interacting* field theory is the first example of a Hodge theory where all the de Rham cohomology operators correspond to some *local* generators.

The outline of this paper is as follows. In Sec. 2, we recapitulate the bare essentials of the BRST symmetries and set up the notations. This is followed by the discussions of dual BRST symmetries in Sec. 3. Section 4 is devoted to the discussion of symmetry generated by the Casimir operator. In Sec. 5, we discuss a discrete symmetry that corresponds to the Hodge duality operation. The extended BRST algebra and the Hodge decomposition theorem are discussed in Sec. 6. Finally, we make some concluding remarks in Sec. 7.

2 BRST Symmetries

Let us begin with a D -dimensional BRST invariant Lagrangian density ($\mathcal{L}_B$) for the interacting $U(1)$ gauge theory in the Feynman gauge:

$$\mathcal{L}_B = -\frac{1}{4}F^{\mu\nu}F_{\mu\nu} + \bar{\psi}(i\gamma^\mu\partial_\mu - m)\psi - e\bar{\psi}\gamma^\mu A_\mu\psi + B(\partial \cdot A) + \frac{1}{2}B^2 - i\partial_\mu\bar{C}\partial^\mu C, \quad (2.1)$$

where $F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu$ is the field strength tensor, B is the Nakanishi-Lautrup auxiliary field, $(\bar{C})C$ are the anticommuting Faddeev-Popov (anti)ghost fields ($\bar{C}^2 = C^2 = 0, C\psi = -\psi C, \bar{C}\psi = -\psi\bar{C}, C\bar{C} = -\bar{C}C$, etc.) and indices $\mu, \nu = 0, 1, 2, \dots, D-1$ represent the flat Minkowski space-time directions. It can be checked that the above Lagrangian density remains quasi-invariant ($\delta_B\mathcal{L}_B = \eta\partial_\mu[B\partial^\mu C]$) under the following off-shell nilpotent ($\delta_B^2 =$

[†]The vector potential A_μ of the $U(1)$ gauge theory is defined through one-form $A = A_\mu dx^\mu$. The gauge-fixing term $\partial \cdot A = \delta A$ is Hodge dual to the two-form $F = dA$ where $\delta = \pm * d*$ is the adjoint(dual) exterior derivative and d is the exterior derivative (see, e.g., Refs. [9,10]).

0) BRST transformations:

$$\begin{aligned}\delta_B A_\mu &= \eta \partial_\mu C, & \delta_B C &= 0, & \delta_B \bar{C} &= i \eta B, & \delta_B B &= 0, \\ \delta_B \psi &= -i\eta e C \psi, & \delta_B \bar{\psi} &= i\eta e C \bar{\psi}, & \delta_B (\partial \cdot A) &= \eta \square C, & \delta_B F_{\mu\nu} &= 0,\end{aligned}\quad (2.2)$$

where η is an anticommuting ($\eta C = -C\eta, \eta \bar{C} = -\bar{C}\eta, \eta \psi = -\psi\eta, \eta \bar{\psi} = -\bar{\psi}\eta$) space-time independent transformation parameter. The on-shell ($\square C = 0$) nilpotent ($\delta_b^2 = 0$) BRST transformations can be obtained from (2.1) by inserting the equations of motion for the B field (i.e. $B = -(\partial \cdot A)$) as

$$\begin{aligned}\delta_b A_\mu &= \eta \partial_\mu C, & \delta_b C &= 0, & \delta_b \bar{C} &= -i \eta (\partial \cdot A), \\ \delta_b \psi &= -i\eta e C \psi, & \delta_b \bar{\psi} &= i\eta e C \bar{\psi}, & \delta_b (\partial \cdot A) &= \eta \square C,\end{aligned}\quad (2.3)$$

and the corresponding on-shell nilpotent BRST invariant Lagrangian density is

$$\mathcal{L}_b = -\frac{1}{4} F^{\mu\nu} F_{\mu\nu} + \bar{\psi} (i\gamma^\mu \partial_\mu - m) \psi - e \bar{\psi} \gamma^\mu A_\mu \psi - \frac{1}{2} (\partial \cdot A)^2 - i \partial_\mu \bar{C} \partial^\mu C. \quad (2.4)$$

It is straightforward to check that the generator for the above symmetry transformations is the off(on)-shell nilpotent ($Q_{B,(b)}^2 = 0$) BRST charge

$$\begin{aligned}Q_B &= \int d^{(D-1)}x [\partial_i F^{0i} C + B \dot{C} - e \bar{\psi} \gamma_0 C \psi] \equiv \int d^{(D-1)}x [B \dot{C} - \dot{B} C], \\ Q_b &= \int d^{(D-1)}x [\partial_i F^{0i} C - (\partial \cdot A) \dot{C} - e \bar{\psi} \gamma_0 C \psi] \\ &\equiv \int d^{(D-1)}x [\partial_0 (\partial \cdot A) C - (\partial \cdot A) \dot{C}],\end{aligned}\quad (2.5)$$

where the latter expressions for $Q_{B,b}$ have been obtained by exploiting the equation of motions: $(\partial_\mu F^{\mu\nu} - \partial^\nu B = e \bar{\psi} \gamma^\nu \psi)$ and $(\partial_\mu F^{\mu\nu} + \partial^\nu (\partial \cdot A) = e \bar{\psi} \gamma^\nu \psi)$. The global scale invariance of (2.1) under $C \rightarrow e^{-\lambda} C, \bar{C} \rightarrow e^\lambda \bar{C}, A_\mu \rightarrow A_\mu, B \rightarrow B$ (where λ is a global parameter), leads to the derivation of a conserved ghost charge (Q_g)

$$Q_g = -i \int d^{(D-1)}x [C \dot{\bar{C}} + \bar{C} \dot{C}]. \quad (2.6)$$

Together, these conserved charges satisfy the following algebra

$$\begin{aligned}Q_B^2 &= \frac{1}{2} \{Q_B, Q_B\} = 0, & i[Q_g, Q_{AB}] &= -Q_{AB}, & i[Q_g, Q_B] &= +Q_B, \\ Q_{AB}^2 &= \frac{1}{2} \{Q_{AB}, Q_{AB}\} = 0, & \{Q_B, Q_{AB}\} &= 0,\end{aligned}\quad (2.7)$$

where Q_{AB} is the anti-BRST charge which can be readily obtained from (2.5) by the replacement $:C \rightarrow i\bar{C}$. Note that $C \rightarrow \pm i\bar{C}, \bar{C} \rightarrow \pm iC$ is the discrete symmetry of the ghost action ($I_{F.P.} = -i \int d^D x \partial_\mu \bar{C} \partial^\mu C$) in any arbitrary dimension of spacetime. This is why the anti-BRST charge Q_{AB} and the corresponding nilpotent transformations can be obtained from (2.5) and (2.2) by the replacement: $C \rightarrow \pm i\bar{C}$.

At this stage, it is worth pointing out the fact that the de Rham cohomology operators (d, δ, Δ) obey the following algebra

$$\begin{aligned}d^2 &= 0, & \delta^2 &= 0, & \Delta &= (d + \delta)^2 = d\delta + \delta d, \\ [\Delta, d] &= 0, & [\Delta, \delta] &= 0 & \{d, \delta\} &= \Delta \neq 0.\end{aligned}\quad (2.8)$$

It can be easily seen that if Q_B is the analogue of d , Q_{AB} is *not* the analogue of δ because Q_B and Q_{AB} anticommute[§] with each-other whereas d and δ do *not* anticommute. Thus, in the algebra (2.7), there is no analogue of the Laplacian operator Δ . Furthermore, under transformations generated by Q_B and Q_{AB} , it is the two-form $F = dA$ (derived from the application of d on the one-form A) that remains invariant and not the gauge-fixing term (which is derived from one-form A by application of dual exterior derivative δ). Thus, it is obvious that, at the algebraic- and conceptual level, the analogy is not complete between the BRST algebra (2.7) and the de Rham cohomology algebra (2.8).

3 Dual BRST symmetries

In two $(1+1)$ dimensions of space-time, there exists only one component (i.e. electric field $E = F_{01}$) of the field strength tensor $F_{\mu\nu}$ and there is no magnetic field. Thus, the analogue of the BRST invariant Lagrangian density (2.1) is:

$$\mathcal{L}_B = \frac{1}{2} E^2 + \bar{\psi}(i\gamma^\mu \partial_\mu - m)\psi - e\bar{\psi}\gamma^\mu A_\mu\psi + B(\partial \cdot A) + \frac{1}{2}B^2 - i\partial_\mu \bar{C}\partial^\mu C, \quad (3.1a)$$

which can be recast as:

$$\mathcal{L}_B = \mathcal{B}E - \frac{1}{2} \mathcal{B}^2 + \bar{\psi}(i\gamma^\mu \partial_\mu - m)\psi - e\bar{\psi}\gamma^\mu A_\mu\psi + B(\partial \cdot A) + \frac{1}{2}B^2 - i\partial_\mu \bar{C}\partial^\mu C, \quad (3.1b)$$

by introducing another auxiliary field $\mathcal{B}$. It can be checked that under the following off-shell nilpotent ($\delta_D^2 = 0$) dual-BRST transformations

$$\begin{aligned} \delta_D A_\mu &= -\eta \varepsilon_{\mu\nu} \partial^\nu \bar{C}, & \delta_D \bar{C} &= 0, & \delta_D C &= -i\eta \mathcal{B}, & \delta_D \mathcal{B} &= 0, & \delta_D B &= 0, \\ \delta_D \psi &= -i\eta e \bar{C} \gamma_5 \psi, & \delta_D \bar{\psi} &= i\eta e \bar{C} \gamma_5 \bar{\psi}, & \delta_D (\partial \cdot A) &= 0, & \delta_D E &= \eta \square \bar{C}, \end{aligned} \quad (3.2)$$

the Lagrangian density (3.1b) (with $m = 0$) transforms as: $\delta_D \mathcal{L}_B = \eta \partial_\mu (\mathcal{B} \partial^\mu \bar{C})$ [¶]. The on-shell ($\square \bar{C} = 0$) nilpotent ($\delta_d^2 = 0$) dual-BRST symmetry transformations (for $m = 0$) can be obtained from (3.2) by exploiting equations of motions w.r.t. B and $\mathcal{B}$ fields (i.e. $B = -(\partial \cdot A)$; $\mathcal{B} = E$) as

$$\begin{aligned} \delta_d A_\mu &= -\eta \varepsilon_{\mu\nu} \partial^\nu \bar{C}, & \delta_d \bar{C} &= 0, & \delta_d C &= -i\eta E, & \delta_d E &= \eta \square \bar{C}, \\ \delta_d \psi &= -i\eta e \bar{C} \gamma_5 \psi, & \delta_d \bar{\psi} &= i\eta e \bar{C} \gamma_5 \bar{\psi}, & \delta_d (\partial \cdot A) &= 0. \end{aligned} \quad (3.3)$$

The above on-shell ($\square \bar{C} = 0$) nilpotent transformations turn out to be the symmetry for the analogue of the Lagrangian density (2.4)

$$\mathcal{L}_b = \frac{1}{2} E^2 + \bar{\psi}(i\gamma^\mu \partial_\mu - m)\psi - e\bar{\psi}\gamma^\mu A_\mu\psi - \frac{1}{2}(\partial \cdot A)^2 - i\partial_\mu \bar{C}\partial^\mu C. \quad (3.4)$$

[§]Cohomologically higher-order BRST- and anti-BRST operators do not anticommute for the compact non-Abelian Lie algebras [23]. Our discussion, however, concerns with a simple Abelian Lie algebra.

[¶]We adopt the notations in which the flat 2D Minkowski metric $\eta_{\mu\nu} = \text{diag}(+1, -1)$, $\gamma^0 = \sigma_2$, $\gamma^1 = i\sigma_1$, $\gamma_5 = \gamma^0 \gamma^1 = \sigma_3$, $\{\gamma^\mu, \gamma^\nu\} = 2\eta^{\mu\nu}$, $\gamma_\mu \gamma_5 = \varepsilon_{\mu\nu} \gamma^\nu$, $\varepsilon_{01} = \varepsilon^{10} = +1$, $F_{01} = \partial_0 A_1 - \partial_1 A_0 = E = -\varepsilon^{\mu\nu} \partial_\mu A_\nu = F^{10}$, $\square = \eta^{\mu\nu} \partial_\mu \partial_\nu = (\partial_0)^2 - (\partial_1)^2$ and here σ 's are the usual 2×2 Pauli matrices.

We christen the above off(on)-shell nilpotent symmetries (3.2) and (3.3) as dual-BRST symmetries because it is the gauge-fixing term that remains invariant under these transformations and the corresponding symmetry transformations on the Dirac fields are the analogue of the chiral transformations. It should be contrasted with the usual BRST symmetries, under which, it is the electric field E that remains invariant and the corresponding symmetry transformations on the Dirac fields are the analogue of the local gauge transformations ^{||}.

It can be checked that the above transformations are generated by

$$\begin{aligned} Q_D &= \int dx [\mathcal{B}\dot{\bar{C}} + e\bar{\psi}\gamma_1\bar{C}\psi - (\partial_1 B)\bar{C}] \equiv \int dx [\mathcal{B}\dot{\bar{C}} - \dot{\mathcal{B}}\bar{C}], \\ Q_d &= \int dx [E\dot{\bar{C}} + e\bar{\psi}\gamma_1\bar{C}\psi + (\partial_1(\partial \cdot A))\bar{C}] \equiv \int dx [E\dot{\bar{C}} - \dot{E}\bar{C}], \end{aligned} \quad (3.5)$$

where the latter expressions for $Q_{D,d}$ have been obtained by using the equation of motion $(\varepsilon^{\mu\nu}\partial_\nu\mathcal{B} + \partial^\mu B = -e\bar{\psi}\gamma^\mu\psi)$ and $(\varepsilon^{\mu\nu}\partial_\nu E - \partial^\mu(\partial \cdot A) = -e\bar{\psi}\gamma^\mu\psi)$ for the photon field present in the Lagrangian density (3.1b) and (3.4). Using the following BRST quantization conditions (with $\hbar = c = 1$):

$$\begin{aligned} [A_0(x, t), B(y, t)] &= i\delta(x - y), [A_1(x, t), \mathcal{B}(y, t)] = i\delta(x - y), \\ \{\psi(x, t), \psi^\dagger(y, t)\} &= -\delta(x - y), \{C(x, t), \bar{C}(y, t)\} = \delta(x - y), \\ \{\bar{C}(x, t), \dot{C}(y, t)\} &= -\delta(x - y), \end{aligned} \quad (3.6)$$

(and rest of the (anti)commutators are zero), it can be seen that Q_D is indeed the generator for the transformations (3.2) ^{**} if we exploit the following relationship

$$\delta_D\Phi = -i\eta[\Phi, Q_D]_\pm, \quad (3.7)$$

where $[\ , \]_\pm$ stands for (anti)commutator for the generic field Φ being (fermionic)bosonic in nature. It is straightforward to check that $Q_D^2 = \frac{1}{2}\{Q_D, Q_D\} = 0$ due to (3.6). A simpler way to see this fact is: $\delta_D Q_D = -i\eta\{Q_D, Q_D\} = 0$ by exploiting (3.2) and (3.5).

4 Symmetry generated by the Casimir operator

It is very natural to expect that the anticommutator of these two transformations $(\{\delta_B, \delta_D\} = \delta_W)$ would also be the symmetry transformation (δ_W) for the Lagrangian density (3.1b) (with $m = 0$). This is indeed the case as can be seen that under the following bosonic $(\kappa = -i\eta\eta')$ transformations corresponding to δ_W

$$\begin{aligned} \delta_W A_\mu &= \kappa(\partial_\mu\mathcal{B} + \varepsilon_{\mu\nu}\partial^\nu B), & \delta_W\mathcal{B} &= 0, & \delta_W B &= 0, \\ \delta_W(\partial \cdot A) &= \kappa\Box\mathcal{B}, & \delta_W E &= -\kappa\Box B, & \delta_W C &= 0, & \delta_W\bar{C} &= 0, \\ \delta_W\psi &= \kappa ie(\gamma_5 B - \mathcal{B})\psi, & \delta_W\bar{\psi} &= -\kappa ie(\gamma_5 B - \mathcal{B})\bar{\psi}, \end{aligned} \quad (4.1)$$

^{||}As explained in Sec. 1, in two dimensional space-time, the electric field E and gauge-fixing term $(\partial \cdot A)$ are ‘Hodge dual’ to each-other. The gauge symmetry and chiral symmetry transformations on the Dirac fields are also dual to each-other (see, e.g., Ref. [24]).

^{**} Here, and in what follows, we shall concentrate only on the off-shell nilpotent symmetries and corresponding generators. The case of on-shell nilpotent symmetries can be easily derived from here.

the Lagrangian density (3.1b) (with $m = 0$) transforms as: $\delta_W \mathcal{L}_B = \kappa \partial_\mu [B \partial^\mu \mathcal{B} - \mathcal{B} \partial^\mu B]$. Here η and η' are the transformation parameters corresponding to the transformations δ_B and δ_D respectively. The generator for the above transformations is:

$$W = \int dx [B(\partial_1 B + e\bar{\psi}\gamma_1\psi) - \mathcal{B}(\partial_1 \mathcal{B} - e\bar{\psi}\gamma_0\psi)] \equiv \int dx [B\dot{\mathcal{B}} - \dot{B}\mathcal{B}], \quad (4.2)$$

where the latter expression for W has been obtained due to the use of equation of motion: $\varepsilon^{\mu\nu}\partial_\nu \mathcal{B} + \partial^\mu B = -e\bar{\psi}\gamma^\mu\psi$. As we shall see below (Sec. 6), this bosonic symmetry generator is the Casimir operator for the full extended BRST algebra which plays an important role in the representation theory for this algebra [11-13]. There are other simpler ways to derive the expression for W . For instance, it can be seen that anticommutator of Q_B and Q_D leads to the derivation of W (i.e. $\{Q_B, Q_D\} = W$) if we exploit the basic (anti)commutators (3.6). Furthermore, since Q_B and Q_D are the generators for transformations (2.2) and (3.2) respectively, it can be seen that the following relationships

$$\begin{aligned} \delta_D Q_B &= -i\eta\{Q_B, Q_D\} = -i\eta W, \\ \delta_B Q_D &= -i\eta'\{Q_D, Q_B\} = -i\eta' W, \end{aligned} \quad (4.3)$$

lead to the definition and derivation of W .

5 Discrete symmetries

It can be readily seen that, in 2D, the ghost action ($I_{F,P} = -i \int d^2x \partial_\mu \bar{C} \partial^\mu C$) remains invariant under

$$C \rightarrow \pm i\bar{C}, \quad \bar{C} \rightarrow \pm iC, \quad \partial_\mu \rightarrow \pm i\varepsilon_{\mu\nu}\partial^\nu. \quad (5.1)$$

This symmetry was exploited for the definition of anti-BRST and dual/anti-dual BRST symmetries in the case of *free* Abelian gauge theories [19, 20]. Furthermore, it was noticed that under the following separate and independent transformations

$$C \rightarrow \pm i\bar{C}, \quad \bar{C} \rightarrow \pm iC \quad \partial_\mu \rightarrow \pm i\varepsilon_{\mu\nu}\partial^\nu \quad A_\mu \rightarrow A_\mu, \quad (5.2)$$

$$C \rightarrow \pm i\bar{C}, \quad \bar{C} \rightarrow \pm iC \quad A_\mu \rightarrow \mp i\varepsilon_{\mu\nu}A^\nu, \quad (5.3)$$

the free (without matter fields ψ and $\bar{\psi}$) Lagrangian density (2.4) remains invariant. The above transformations were found to be the analogue of Hodge duality (*) operations for the free theory [19,20]. It was also shown that the (anti)dual BRST transformations can be obtained from the usual (anti)BRST transformations by exploiting the above discrete symmetries. Similarly, two sets of topological invariants of the theory were shown to be connected to each-other by (5.2) or (5.3).

For the interacting theory, described by the Lagrangian densities (3.1a, 3.1b) and (3.4), similar transformations exist. It can be checked that, under the following transformations

$$\begin{aligned} C &\rightarrow \pm i\gamma_5 \bar{C}, & A_0 &\rightarrow \pm i\gamma_5 A_1, & E &\rightarrow \pm i\gamma_5 (\partial \cdot A), & \psi &\rightarrow \psi, \\ \bar{C} &\rightarrow \pm i\gamma_5 C, & A_1 &\rightarrow \pm i\gamma_5 A_0, & (\partial \cdot A) &\rightarrow \pm i\gamma_5 E, & \bar{\psi} &\rightarrow \bar{\psi}, \\ \mathcal{B} &\rightarrow \mp i\gamma_5 B, & B &\rightarrow \mp i\gamma_5 \mathcal{B}, & e &\rightarrow \mp ie, \end{aligned} \quad (5.4)$$

the above Lagrangian densities (for the interacting theory) remain invariant ^{††}. Furthermore, it can be easily seen that the dual-BRST symmetries (3.2) and (3.3) can be derived from their counterpart BRST symmetries ((2.2) and (2.3)) by exploiting the symmetry transformations (5.4). Thus, as argued in Refs. [19, 22], the discrete transformations in (5.4) are the analogue of the Hodge duality (*) operation of differential geometry. To verify this claim, we first note that, under two successive * operations (corresponding to transformations (5.4)), the generic fields Φ of the above Lagrangian densities, transform as

$$* (* \Phi) = \pm \Phi, \quad (5.5)$$

where (+) sign stands for the generic field Φ being $\psi, \bar{\psi}$ and (-) sign is for the rest of the fields (e.g. $\Phi = C, \bar{C}, B, \mathcal{B}, A_0, A_1, (\partial \cdot A), E$). Now, it is very gratifying to note that, for the generic field Φ , we have the following relationship

$$\delta_D \Phi = \pm * \delta_B * \Phi, \quad (5.6)$$

where δ_D and δ_B are the nilpotent symmetry transformations (3.2) and (2.2) respectively and (+) sign stands for $\Phi = \psi, \bar{\psi}$ and (-) sign is for the rest of the fields. The relative signs in (5.6) are dictated by the corresponding signs in (5.5). Thus, we see that the relationship between transformations δ_B and δ_D acting on the generic field Φ is same as the relationship between exterior derivative d and the dual-exterior derivative δ (i.e. $\delta = \pm * d *$) acting on a differential form (defined on a compact manifold).

It is worthwhile to mention that under discrete transformations (5.4), the generators ($Q_{(B,AB)}, Q_{(D,AD)}, W$) for the continuous transformations undergo the following change

$$Q_{(B,AB)} \rightarrow Q_{(D,AD)}, \quad Q_{(D,AD)} \rightarrow Q_{(B,AB)} \quad W \rightarrow W, \quad (5.7)$$

so that the extended BRST algebra (see, e.g., eqn. (6.1) below) remains intact as far as these operators are concerned. Here Q_{AD} is the anti-dual BRST charge which can be readily derived from (3.5) by the replacement $:\bar{C} \rightarrow \pm i C$. The above transformations (5.7) should be contrasted with the four (3+1) dimensional case of the free 2-form Abelian gauge theories [22] where it has been shown that under the analogue of * operation, the generators transform as: $Q_B \rightarrow Q_D, Q_D \rightarrow -Q_B, W \rightarrow -W$ which turns out to mimic the duality symmetry transformation of the free Maxwell equations where $\mathbf{E} \rightarrow \mathbf{B}, \mathbf{B} \rightarrow -\mathbf{E}$. As expected by the duality considerations in 2D [25-27], the Hodge duality (*) transformations in (5.7) are correct and consistent because of the fact that dualities in 4D and 2D are of different nature. In fact, it is precisely because of this reason that, in 2D, one obtains (a reciprocal relationship between δ_B and δ_D):

$$\delta_B \Phi = \pm * \delta_D * \Phi, \quad (5.8)$$

^{††}Note that in the matrix notations, the transformations like: $A_0 \rightarrow \pm i \gamma_5 A_1$, etc., imply that $A_0 \rightarrow \pm i A_1$ or/and $A_0 \rightarrow \mp i A_1$. Except for the exchange of signs, these transformations are same. In fact, γ_5 is present appropriately in the rest of the transformations to take care of this sign-flip.

analogous to (5.6). This is a reflection of the fact that in 2D, there is no relative change of sign between an operator and its dual operator under the duality transformation.

6 Extended BRST algebra

Together, all the above generators obey the following extended BRST algebra

$$\begin{aligned}
[W, Q_k] &= 0, & k &= g, B, D, AB, AD, \\
Q_B^2 &= Q_D^2 = Q_{AB}^2 = Q_{AD}^2 = 0, & \{Q_D, Q_{AD}\} &= 0, \\
\{Q_B, Q_D\} &= \{Q_{AB}, Q_{AD}\} = W, & \{Q_B, Q_{AB}\} &= 0, \\
i[Q_g, Q_B] &= Q_B, & i[Q_g, Q_{AB}] &= -Q_{AB}, \\
i[Q_g, Q_D] &= -Q_D, & i[Q_g, Q_{AD}] &= Q_{AD},
\end{aligned} \tag{6.1}$$

and the rest of the (anti)commutators are zero. It is evident that the operator W is the Casimir operator for the whole algebra. The *mathematical aspects* of the representation theory for the above kind of BRST algebra have been discussed in Refs. [11-13]. It will be noticed that the ghost number for Q_B and Q_{AD} is +1 and that of Q_D and Q_{AB} is -1. Now, given a state $|\phi\rangle$ with ghost number n in the quantum Hilbert space (i.e. $iQ_g|\phi\rangle = n|\phi\rangle$), it can be readily seen, using the algebra (6.1), that

$$\begin{aligned}
iQ_g Q_B |\phi\rangle &= (n+1)Q_B |\phi\rangle, & iQ_g Q_{AD} |\phi\rangle &= (n+1)Q_{AD} |\phi\rangle, \\
iQ_g Q_D |\phi\rangle &= (n-1)Q_D |\phi\rangle, & iQ_g Q_{AB} |\phi\rangle &= (n-1)Q_{AB} |\phi\rangle, \\
iQ_g W |\phi\rangle &= n W |\phi\rangle.
\end{aligned} \tag{6.2}$$

This shows that the ghost numbers for the states $Q_B|\phi\rangle$ (or $Q_{AD}|\phi\rangle$), $Q_D|\phi\rangle$ (or $Q_{AB}|\phi\rangle$) and $W|\phi\rangle$ in the quantum Hilbert space are $(n+1)$, $(n-1)$ and n respectively.

At this stage, it is worth mentioning that in Refs. [19-21], the analogous expressions for Q_B, Q_D, W have been derived for the free 2D Abelian- and non-Abelian gauge theories (having no interaction with matter fields). The topological properties of these free theories have been shown to be encoded in the vanishing of the operator W when equations of motion are exploited. On the contrary, as it turns out, the operator W is defined off-shell as well as on-shell for the 2D interacting BRST invariant $U(1)$ gauge theory. This is because of the fact that even though $U(1)$ gauge field is topological (i.e. without any propagating degrees of freedom), it is coupled to the Dirac fields here and fermionic degrees of freedom are present in the off-shell as well as on-shell expression for W . Thus, the present theory is an example of an *interacting* topological field theory in 2D.

It is interesting to note that the algebra of Q_B, Q_D and W in equation (6.1) is exactly identical to the corresponding algebra for the de Rham cohomology operators (d, δ, Δ) in (2.8). Furthermore, it can be readily seen that the operation of these generators on a state with ghost number n (cf. eqn.(6.2)) is same as the operation of the above cohomological operators on a differential form of degree n . Thus, it is clear that the BRST cohomology can be defined comprehensively in terms of the above operators and Hodge decomposition theorem can be implemented cogently in the quantum Hilbert space of states where any

arbitrary state $|\phi\rangle_n$ (with ghost number n) can be written as the sum of a harmonic state $|\omega\rangle_n$ ($W|\omega\rangle_n=0, Q_B|\omega\rangle_n=0, Q_D|\omega\rangle_n=0$), a BRST exact state ($Q_B|\theta\rangle_{n-1}$) and a co-BRST exact state ($Q_D|\chi\rangle_{n+1}$). Mathematically, this statement (which is the analogue of eqn. (1.1)) can be expressed by the following equation

$$|\phi\rangle_n = |\omega\rangle_n + Q_B|\theta\rangle_{n-1} + Q_D|\chi\rangle_{n+1} . \quad (6.3)$$

It is obvious, therefore, that the above symmetry generators Q_B, Q_D and W have their counterparts in differential geometry as the de Rham Cohomology operators d, δ and Δ respectively [9,10]. It is an interesting feature of the BRST formalism that these de Rham cohomological operators can also be identified with the generators Q_{AD}, Q_{AB} and $W = \{Q_{AD}, Q_{AB}\}$ respectively. Thus, there is a two-to-one mapping from BRST formalism to the differential geometry as: $(Q_B, Q_{AD}) \Rightarrow d, (Q_D, Q_{AB}) \Rightarrow \delta, W = \{Q_B, Q_D\} = \{Q_{AD}, Q_{AB}\} \Rightarrow \Delta$.

7 Conclusions

We have shown that, in the language of well-defined symmetry properties of the BRST invariant Lagrangian density of an interacting $U(1)$ gauge theory, the celebrated Hodge decomposition theorem can be understood. The generators of the nilpotent BRST transformations; nilpotent dual BRST transformations and a bosonic symmetry transformation (derived by taking the anticommutator of both these nilpotent symmetries) correspond to the de Rham cohomology operators (d, δ, Δ) of differential geometry and they obey the same kind of algebra as obeyed by these operators. There is a logical understanding for the existence of these generators. In fact, these individual generators generate *separate and independent* symmetry transformations under which the two-form field (i.e. electric field E), gauge-fixing term and ghost fields remain invariant respectively. A discrete symmetry of the theory corresponds to the Hodge duality $(*)$ operation of the differential geometry. Thus, all the cohomological quantities of differential geometry find their analogue in the language of physically well-defined quantities for interacting $U(1)$ gauge theory. Thus, this interacting theory is a physical field theoretical model for the Hodge theory. It will be very useful to explore the impact of this new symmetry in the context of symmetries of the Green's functions for QED and derive the analogue of Ward-Takahashi identities. Since the new symmetry is connected with the analogue of chiral transformation, it might be possible that BRST cohomology and Hodge decomposition theorem will shed some light on the Adler-Bardeen-Jackiw anomaly in 2D. It is quite possible that this understanding might provide an insight into the proof of the consistency and unitarity of the anomalous gauge theory in two-dimensions (see, e.g., Refs. [28,29] and references therein). The generalization of this new symmetry to 2D non-Abelian gauge theory (having local gauge interaction with matter fields) is another future direction that can be pursued. A very good attempt has been made to discuss these new symmetries in the framework of Hamiltonian formulation [30]. However, more works are needed in this direction to establish a precise analogy with the key concepts of differential geometry. The insights gained in these studies might

provide a clue for the generalization of this new symmetry to physical four dimensional gauge theories. These are some of the issues under investigation and a detailed discussion will be reported elsewhere.

Acknowledgments

A part of this work was done at JINR, Dubna (Moscow) and AS-ICTP, Trieste (Italy). Fruitful conversations with members of theory groups at BLTP (JINR) and HEP group at AS-ICTP are gratefully acknowledged.

References

- [1] C. Becchi, A. Rouet and R. Stora, Phys. Lett. **B32** (1974) 344; Commun. Math. Phys. **42** (1975) 127; Ann. Phys. (N.Y.) **98** (1976) 28.
- [2] I. V. Tyutin, *Lebedev Institute Report No.* FIAN-39 (1975) (unpublished).
- [3] P. A. M. Dirac, *Lectures on Quantum Mechanics*, (Yeshiva University Press, New York, 1964).
- [4] K. Sundermeyer, *Constrained Dynamics: Lecture Notes in Physics* (Springer, New York, 1982).
- [5] D. M. Gitman and I. V. Tyutin, *Quantization of Fields with Constraints*, (Springer, Berlin, 1990).
- [6] K. Nishijima, in: *Progress in Quantum Field Theory* eds. H. Ezawa and S. Kamefuchi (North-Holland, Amsterdam, 1986) p. 99.
- [7] M. Henneaux and C. Teitelboim, *Quantization of Gauge Systems* (Princeton University Press, Princeton NJ, 1992).
- [8] N. Nakanishi and I. Ojima, *Covariant Operator Formalism of Gauge Theories and Quantum Gravity* (World Scientific, Singapore, 1990).
- [9] T. Eguchi, P. B. Gilkey and A. J. Hanson, Phys. Rep. **66** (1980) 213.
- [10] S. Mukhi and N. Mukunda, *Introduction to Topology, Differential Geometry and Group Theory for Physicists* (Wiley Eastren Ltd., New Delhi, 1990).
- [11] K. Nishijima, Prog. Theor. Phys. **80** (1988) 897; 905.
- [12] J. W. van Holten, Phys. Rev. Lett. **64** (1990) 2863; Nucl. Phys. **B339** (1990) 158.
- [13] G. Fülöp and R. Marnelius, Nucl. Phys. **B456** (1995) 442.
- [14] D. McMullan and M. Lavelle, Phys. Rev. Lett. **71** (1993) 3758; *ibid.* **75** (1995) 4151.
- [15] V. O. Rivelles, Phys. Rev. Lett. **75** (1995) 4150, Phys. Rev. **D 53** (1996) 3257.
- [16] R. Marnelius, Nucl. Phys. **B494** (1997) 346.
- [17] H. S. Yang and B. -H. Lee, J. Math. Phys. **37** (1996) 6106.

- [18] T. Zhong and D. Finkelstein, Phys. Rev. Lett. **73** (1994) 3055; *ibid.* **75** (1995) 4152.
- [19] R. P. Malik, Int. J. Mod. Phys. **A15** (2000) 1685 : hep-th/9808040; J. Phys. A: Math Gen **33** (2000) 2437: hep-th/9902146.
- [20] R. P. Malik, Mod. Phys. Lett. **A14** (1999) 1937: hep-th/9903128.
- [21] R. P. Malik, *New topological field theories in two dimensions* : hep-th/0012085.
- [22] E. Harikumar, R. P. Malik and M. Sivakumar, J. Phys. A: Math Gen **33** (2000) 7149: hep-th/0004145.
- [23] C. Chryssomalakos, J. A. de Azcarraga, A. J. Macfarlane, and J. C. Perez Bueno, J. Math. Phys. **40** (1999) 6009.
- [24] R. P. Malik and T. Pradhan, Z. Phys. **C 28** (1985) 525.
- [25] S. Deser, A. Gomberoff, M. Henneaux and C. Teitelboim, Phys. Lett. **B 400** (1997) 80.
- [26] R. Banerjee and B. Chakraborty, J. Phys. A: Math Gen **32** (1999) 4441.
- [27] R. Banerjee and C. Wotzasek, Nucl. Phys. **B 527** (1998) 402.
- [28] R. Jackiw and R. Rajaraman, Phys. Rev. Lett. **54** (1985) 1219.
- [29] R. P. Malik, Phys. Lett. **B212** (1988) 445.
- [30] P. Gaete, *Note on the dual BRST symmetry in $U(1)$ gauge theory*: hep-th/0005203.